

Mesothelioma mortality by occupation statistics in Great Britain, 2023



July 2023



Contents

Summary	2
Introduction	2
Results	4
Data tables	4
Methods and limitations	4
Overall PMRs for 2011-2021 and time trends for 2001-2021	6
References	19
Annex 1 – Technical notes	20
Example PMR calculation	21
Confidence intervals and statistical significance	22
National Statistics	23

Summary

The information in this document relates to Health and Safety Statistics published by the Health and Safety Executive in 2023. The document can be found at:

www.hse.gov.uk/statistics/causdis

Introduction

This fact sheet presents updated mesothelioma mortality statistics by last recorded occupation of the deceased within Great Britain from 2011-2021 and time trends over the longer period of 2001-2021. The document can be found at

www.hse.gov.uk/statistics/causdis/index.htm.

Background information about mesothelioma and statistics for mesothelioma deaths in Great Britain as a whole and by geographical area within Great Britain are also available at

www.hse.gov.uk/statistics/causdis/index.htm.

These statistics are based on the last occupation of the deceased, as recorded on death certificates. The Proportional Mortality Ratio (PMR) presented for each occupation compares the frequency that the occupation is recorded for mesothelioma deaths with the frequency that it is recorded for deaths from all causes of death as a whole.

PMRs thus provide a way of highlighting occupations that may be associated with higher-than-average mortality from mesothelioma.

The analyses of temporal trends in occupational PMRs within Great Britain should be interpreted as an indication of how the proportion of deaths with a particular occupation recorded has changed over time, rather than the absolute numbers.

The PMR statistics are limited by the fact that only the last occupation of the deceased is recorded on death certificates which, given the typically long period that the disease takes to develop, may not always be the relevant occupation in terms of past exposure to asbestos. The analysis is restricted to deaths occurring at ages 16-74 years since occupations are routinely recorded on death certificates only for deaths in this range in England and Wales.

Overall deaths increased more than 35% over the period of analysis (i.e. since 2001), and around 10-fold since the late 1960s when consistent recording of mesothelioma in Britain began. However, deaths below age 75 years – those included in this analysis – have reduced over the last 10 years. Nevertheless, since the value of these statistics is in the relative comparison of the frequency of recording of different occupations on mesothelioma death certificates rather than in the absolute number of deaths for each occupation, the statistics still provide useful information about the occupations are more likely to have been a source of past asbestos exposure.




Figure 1: Male and female mesothelioma deaths 2001-2021 by age category

Results

Data tables

Full results of the PMR calculations by occupation in Great Britain are available in Excel tables at:

www.hse.gov.uk/statistics/tables/mesooccupation.xlsx.

Tables show the number of mesothelioma deaths and PMRs for both males and females by Standard Occupational Classification (SOC) major (1-digit code), sub-major (2-digit code), minor (3-digit code) and unit (4-digit code) groups based on:

- MESO OCCUPATION1: 2011-20 SOC2010 male
- MESO OCCUPATION1a: 2011-21 SOC2010 male
- MESO OCCUPATION2: 2011-20 SOC2010 female
- MESO OCCUPATION2a: 2011-21 SOC2010 female
- MESO OCCUPATION3: 2001-10 SOC2010 male
- MESO OCCUPATION4: 2001-10 SOC2010 female

Note that coding by SOC2020 is not yet generally available for deaths in 2021. Results have been published for the SOC2010 period of 2011-2020 and for an extended period 2011-2021.

SOC codes form a nested hierarchy: the first digit of any full 4-digit unit group code gives its major group, the first two digits gives it sub-major group and the first three digits gives its minor group.

Tables include ranks from highest to lowest PMR within each 1- to 4-digit level separately (groups with 10 or fewer observed or expected mesothelioma deaths are not included in the rankings due to the uncertainty associated with smaller numbers).

Methods and limitations

The observed number of deaths in a particular occupation does not represent the actual number of deaths that are attributable to asbestos exposures in that occupation.

PMRs summarise mortality among occupational groups relative to the average level for all occupations in Great Britain as a whole and do not represent absolute measures of risk.

PMRs are expressed as a percentage: values higher or lower than 100 indicate mesothelioma rates that are higher or lower, respectively, than the average for all occupations combined. The corresponding confidence interval should be used to assess whether such an effect could merely be due to random variation.

Occupations with the highest PMRs and where the lower limit of the associated Confidence Interval (CI) are above 100 constitute those that can most reliably be said to have an excess of mesothelioma deaths compared to the average for all occupations, and are, therefore, those most likely to be reflecting an effect due to past occupational asbestos exposure.

Last occupation of the deceased

These analyses are limited by the fact that death certificates record only the last occupation of the deceased. For example, a case of mesothelioma caused by work in the construction industry will only be assigned to that occupation in this analysis if the individual is still in that kind of work when they retired (or died). The long latency period of mesothelioma means that individuals may move between occupations before the onset of the disease and thus there is considerable potential for dilution of the observed difference in risk between occupations.

The dilution will be stronger for those kinds of work where there have been substantial reductions in the relevant workforce (e.g. shipyards, railway rolling stock). The occupations with the highest PMRs will tend to be those which are genuine sources of risk, but PMRs may understate the true relative risk level. PMRs of other occupations will overstate the level of risk (if any) associated with these jobs; occupations with the lowest PMRs will be those which do not entail asbestos exposure, and which are unlikely to be the final full-time occupation for individuals with asbestos exposure.

Role of environmental asbestos exposure

Occupation is recorded on death certificates for deaths at ages 16-74 as a matter of course: for mesothelioma deaths occupation is recorded regardless of whether the deaths were caused by 'occupational exposure' to asbestos. This is particularly important to the interpretation of mesothelioma PMRs for women. Whilst some occupations are recorded as the last occupation on female mesothelioma deaths in appreciable numbers, fewer occupations show evidence that the PMRs are increased. Those occupations that do show increased PMRs in women are not those where the direct handling of asbestos materials at work was likely to have been taking place routinely. Many of these deaths may reflect 'environmental' asbestos exposure, which potentially included any exposures accrued indirectly in the working environment.

Deaths occurring in the latest 10/11-year period (2011-20 and 2011-2021) still predominantly relate to the cohort of people who were younger during the period of peak asbestos use in the 1960s and 1970s when there were far less stringent controls that

required today (e.g. see tables MESOOCUPATION01 and MESOOCUPATION02 which show results for 2011-20 deaths by birth cohort).

The latest occupational analyses of female mesothelioma deaths suggest there is some variation in the average risk of mesothelioma among those who worked in jobs not involving the use of asbestos. For example, proportional mortality ratios are somewhat higher for teachers and administrative occupations than those for nurses, sales occupations and process operatives, and this may suggest the potential for asbestos exposure during work time was somewhat higher in these jobs during the period of peak use. However, past exposures in buildings may have contributed to the background risk seen across all of these kinds of jobs to some extent, and other sources of exposure – for example, in housing stock – are also likely to have contributed.

Other research confirms that, while still caused by asbestos, a majority of mesotheliomas among women (and a similar absolute number among men, though these constitute a smaller proportion of the larger male total) were not directly attributable to occupational or domestic asbestos exposures [note 1]. This, together with an overall increase in mesothelioma deaths among women, suggests there was an increase in the ‘background’ risk among those who did not work with asbestos, but who lived through the period of peak asbestos use. During this period the opportunities for unwitting exposure may have been widespread. This background risk – which has since reduced [note 2] – is likely to at least partly account for deaths with occupations not typically associated with asbestos exposure recorded on the death certificate. The background risk will also apply to men of the same generation.

Reliability of unit group coding

The coding of occupation is likely to be more reliable at the minor group (3-digit code) level than the unit group (4-digit code) level since the recorded information about the job title on death certificates does not always give sufficient information to accurately assign a 4-digit code.

Overall PMRs for 2011-2021 and time trends for 2001-2021

This section presents time trends in PMRs for selected occupations within different levels of the SOC hierarchy where occupational categories based on SOC2000 and SOC2010 were equivalent.

Trends for a particular occupation indicate how the proportion of deaths with a particular occupation recorded has changed over time, rather than the absolute numbers.

The charts show trend lines with solid bold **black** lines to indicate a statistically significant annual trend. Those with **green** lines indicate trends of borderline significance, and for those with **blue** lines trends were not significant. The dashed lines represent the 95% confidence intervals.




Figure 2: Mesothelioma PMRs by SOC major group, males, 2001-2021

SOC major group (1-digit)

Among males, major group 5 (Skilled trades occupations) was the only major group with statistically significantly elevated mesothelioma mortality (PMR=164.7, 95% CI: 159.9, 169.6), with 4478 deaths amongst those aged 16-74 for the period 2011-21. (2011-2020 PMR=163.7, 95% CI: 158.8, 168.8, with 4162 deaths.) This major group contains a number of more specific codes with significantly elevated PMRs, including the only two elevated 2-digit codes, seven of the ten highest ranking 3-digit codes and the 1st (5315: Carpenters and joiners), 3rd (5314: Plumbers and heating and ventilating engineers), 4th (5216: Pipe fitters), 5th (5236: Boat and ship builders and repairers), 6th (5241: Electricians and electrical fitters), 9th (5225: Air-conditioning and refrigeration engineers) and 10th (5322: Floorers and wall tilers) highest ranking 4-digit codes.

The remaining eight major groups generally have consistently significantly lower PMRs compared to the average for all occupations.

Figure 2 shows the temporal trends in the mesothelioma PMRs for males for the nine SOC major groups. There was little evidence of any change in the PMRs over the period 2001-2021 at the major group level.

Among females there were two major groups with statistically significantly elevated mesothelioma mortality:-

- Major group 4: Administrative and secretarial occupations (438 deaths, PMR=127.8, 95% CI: 116.1, 140.3, and evidence of an increasing trend). (2011-2020: 399 deaths, PMR=124.8, 95% CI: 112.8, 137.6.)
- Major group 2: Professional Occupations (258 deaths, PMR=116.6, 95% CI: 102.8, 131.7). (2011-2020: 241 deaths, PMR=117.3, 95% CI: 103.0, 133.1).

Otherwise, there was no evidence of any change in the PMRs over the period 2001-2021 for these groups (Figure 3).




Figure 3: Mesothelioma PMRs by SOC major group, females, 2001-2021

SOC sub-major group (2-digit)

There were two statistically significantly elevated sub-major occupational groupings in the period 2011-2021 for males:

- Group 53: Skilled construction and building trades (2590 deaths, PMR=246.9, 95% CI: 237.5, 256.6) (2011-2020: 2412 deaths, PMR=245.9, 95% CI: 236.2, 255.9.) ranked 1st.
- Group 52: Skilled metal, electrical and electronic trades (1600 deaths, PMR=146.5, 95% CI: 139.4, 153.9). (2011-2020: 1496 deaths, PMR=146.0, 95% CI: 138.7, 153.6) ranked 10th.

The corresponding PMRs for SOC2000 codes for 2001-2010 were also similarly elevated.

Figures 4 shows the results of the trend analyses for these two sub-major groups. There is some evidence of a reduction in the PMR for sub-major group 52 over time.




Figure 4: Mesothelioma PMRs for SOC sub-major group 52 and 53, males, 2001-2021

For females (see figure 4a), the top three and sixth of the top six sub-major groups were statistically significantly elevated during the period 2011-2021:

- Group 23: Teaching and educational professionals (120 deaths, PMR=140.6, 95% CI: 116.5, 168.1). (2011-2020: 113 deaths, PMR=141.9, 95% CI: 116.9, 170.6). The PMR for the corresponding SOC2000 code for 2001-2010 was not elevated.
- Group 35: Business and public service associate professionals (54 deaths, PMR=135.2, 95% CI: 101.6, 176.5). (2011-2020: 48 deaths, PMR=130.0, 95% CI: 95.9, 172.4). The PMR for the corresponding SOC2000 code for 2001-2010 was not elevated.
- Group 41: Administrative occupations (279 deaths, PMR=133.1, 95% CI: 117.9, 149.7). (2011-2020: 254 deaths, PMR=130.9, 95% CI: 115.2, 148.0). The PMR for the corresponding SOC2000 code for 2001-2010 was not elevated.

- Group 91: Elementary trades and related occupations (57 deaths, PMR=129.9, 95% CI: 98.4, 168.3). (2011-2020: 54 deaths, PMR=131.6, 95% CI: 98.9, 172.8). The PMR for the corresponding SOC2000 code for 2001-2010 was elevated.
- Group 24: Business, Media and public service professionals (40 deaths, PMR=119.5, 95% CI: 85.4, 162.8). (2011-2020: 37 deaths, PMR=120.3, 95% CI: 84.7, 165.8). The PMR for the corresponding SOC2000 code for 2001-2010 was not elevated.
- Group 42: Secretarial and related occupations (159 deaths, PMR=119.3, 95% CI: 101.5, 139.4). (2011-2020: 145 deaths, PMR=115.4, 95% CI: 97.4, 135.8). The PMR for the corresponding SOC2000 code for 2001-2010 was not elevated




Figure 4a: Mesothelioma PMRs for SOC sub-major groups 23, 41 and 91, females, 2001-2021

SOC minor group (3-digit)

For males, mesothelioma PMRs for eight SOC minor groups were statistically significantly elevated for the period 2011-2021, all of which have at least some association with building-related activities:

- Group 531: Construction and Building Trades (2135 deaths, PMR=274.3, 95% CI:262.8, 286.2) ranked 1st.
- Group 524: Electrical and Electronic Trades (747 deaths, PMR=211.2, 95% CI:196.3, 226.9) ranked 2nd.
- Group 532: Building Finishing Trades (424 deaths, PMR=170.5, 95% CI:154.7, 187.6) ranked 3rd.
- Group 814: Construction Operatives (203 deaths, PMR=159.8, 95% CI:138.6, 183.3) ranked 4th.
- Group 521: Metal Forming, Welding and Related Trades (214 deaths, PMR=141, 95% CI:122.7, 161.2) ranked 6th.
- Group 812: Plant and Machine Operatives (417 deaths, PMR=125.6, 95% CI:113.8, 138.2) ranked 11th.
- Group 522: Metal Machining, Fitting and Instrument Making Trades (434 deaths, PMR=125.1, 95% CI:113.6, 137.4) ranked 12th.
- Group 243: Architects, Town Planners and Surveyors (107 deaths, PMR=123, 95% CI:100.8, 148.6) ranked 14th.




Figure 5A: Mesothelioma PMRs for selected SOC minor groups, males, 2001-2021

There is some evidence of a reduction in the PMR for minor group 521 Metal Forming, Welding and Related Trades (Figures 5A & 5B).




Figure 5B: Mesothelioma PMRs for selected SOC minor groups, males, 2001-2021

For females, mesothelioma PMRs for five SOC minor groups were statistically significantly elevated for the period 2011-2021:

- Group 241: Legal Professionals (10 deaths, PMR=274.8, 95% CI:131.9, 505.3) ranked 1st.
- Group 413: Administrative Occupations, Records (34 deaths, PMR=185.7, 95% CI:128.6, 259.5) ranked 2nd.
- Group 921: Elementary Administration Occupations (16 deaths, PMR=170.9, 95% CI:97.7, 277.5) ranked 3rd.
- Group 415: Other Administrative Occupations (103 deaths, PMR=160.4, 95% CI:130.9, 194.6) ranked 4th.
- Group 231: Teaching and Educational Professionals (120 deaths, PMR=140.6, 95% CI:116.5, 168.1) ranked 9th.
- Group 421: Secretarial and Related Occupations (159 deaths, PMR=119.3, 95% CI:101.5, 139.4) ranked 16th.

Figure 5C shows an increasing trend amongst 231 (Teaching and educational professionals).




Figure 5C: Mesothelioma PMRs for selected SOC minor groups, females, 2001-2021

SOC unit group (4-digit)

For males, PMRs were statistically significantly elevated for 24 of the 186 SOC unit groups with at least 10 observed or expected mesothelioma deaths. Results for these groups are listed below. Again, a substantial proportion of these unit groups were associated with building activities.

Unit groups with the highest PMRs (higher than 300 the top four):

- 5315: Carpenters and joiners (1037 deaths, PMR=485.3, 95% CI:456.2, 515.8) 1st
- 8124: Energy plant operatives (38 deaths, PMR=343.7, 95% CI:243.2, 471.8) 2nd
- 5314: Plumbers and heating and ventilating engineers (473 deaths, PMR=342.4, 95% CI:312.2, 374.7) 3rd
- 5216: Pipe fitters (55 deaths, PMR=313.8, 95% CI:236.4, 408.4) 4th

Unit groups with high PMRs (PMR of 200 to 300, 5th to 13th):

- 5236: Boat and ship builders and repairers (69 deaths, PMR=279.6, 95% CI:217.5, 353.8) ranked 5th.

- 5241: Electricians and electrical fitters (609 deaths, PMR=272.5, 95% CI:251.3, 295) ranked 6th .
- 2123: Electrical engineers (17 deaths, PMR=271.4, 95% CI:158.1, 434.6) ranked 7th .
- 2424: Business and financial project management professionals (48 deaths, PMR=269.3, 95% CI:198.5, 357) ranked 8th .
- 5225: Air-conditioning and refrigeration engineers (20 deaths, PMR=248.1, 95% CI:151.6, 383.2) ranked 9th .
- 5322: Floorers and wall tilers (59 deaths, PMR=225.7, 95% CI:171.8, 291.2) ranked 10th .
- 5213: Sheet metal workers (58 deaths, PMR=223.5, 95% CI:169.7, 288.9) ranked 11th .
- 1122: Production managers and directors in construction (117 deaths, PMR=204.3, 95% CI:169, 244.9) ranked 12th .
- 1259: Managers and proprietors in other services n.e.c. (184 deaths, PMR=202.5, 95% CI:174.2, 233.9) ranked 13th .

Other unit groups with elevated PMRs (PMRs of 100 to 200, 14th to 22nd, 28th and 30th):

- 8149: Construction operatives n.e.c. (125 deaths, PMR=195.5, 95% CI:162.7, 233) ranked 14th .
- 8125: Metal working machine operatives (276 deaths, PMR=192.2, 95% CI:170.2, 216.3) ranked 15th .
- 5442: Furniture makers and other craft woodworkers (40 deaths, PMR=185.8, 95% CI:132.8, 253.1) ranked 16th .
- 8141: Scaffolders, staggers and riggers (53 deaths, PMR=180.3, 95% CI:135.1, 235.9) ranked 17th .
- 5319: Construction and building trades n.e.c. (474 deaths, PMR=178.4, 95% CI:162.7, 195.2) ranked 18th .
- 5323: Painters and decorators (309 deaths, PMR=173.8, 95% CI:154.9, 194.3) ranked 19th .
- 3563: Vocational and industrial trainers and instructors (36 deaths, PMR=171.3, 95% CI:120, 237.2) ranked 20th .
- 5214: Metal plate workers, and riveters (22 deaths, PMR=166.4, 95% CI:104.3, 252) ranked 21st .
- 1139: Functional managers and directors n.e.c. (37 deaths, PMR=164.3, 95% CI:115.6, 226.4) ranked 22nd .
- 5223: Metal working production and maintenance fitters (346 deaths, PMR=138.1, 95% CI:124, 153.5) ranked 30th .
- 5249: Electrical and electronic trades n.e.c. (63 deaths, PMR=131.3, 95% CI:100.9, 167.9) ranked 35th .




Figure 6A: Mesothelioma PMRs for selected SOC unit groups, males, 2001-2021




Figure 6B: Mesothelioma PMRs for selected SOC unit groups, males, 2001-2021




Figure 6C: Mesothelioma PMRs for selected SOC unit groups, males, 2001-2021




Figure 6D: Mesothelioma PMRs for selected SOC unit groups, males, 2001-2021

For females, PMRs were statistically significantly elevated for five of the 38 SOC unit groups with at least 10 observed or expected mesothelioma deaths:

- 9219: Elementary administration occupations n.e.c. (13 deaths, PMR=317.2, 95% CI:168.8, 542.4) ranked 1st.
- 4131: Records clerks and assistants (17 deaths, PMR=248.7, 95% CI:144.8, 398.2) ranked 2nd.
- 2315 Primary and nursery education teaching professionals (87 deaths, PMR=211.5, 95% CI:169.4, 260.9) ranked 3rd.
- 4211: Medical secretaries (18 deaths, PMR=176.6, 95% CI:104.7, 279.1) ranked 5th.
- 4159: Other administrative occupations n.e.c. (103 deaths, PMR=165.8, 95% CI:135.3, 201.1) ranked 6th.

References

1. Rake C, Gilham C, Hatch J, et al. Occupational, domestic and environmental mesothelioma risks in the British population: a case control study. *British Journal of Cancer* 2009;100(7):1175-83.
2. Gilham C, Rake C, Hodgson J at al. Past and current asbestos exposure and future mesothelioma risks in Britain: The Inhaled Particles Study (TIPS). *International Journal of Epidemiology* 2018;47(6):1745-1756.

Annex 1 – Technical notes

These analyses are based on 49% of male and 41% of female mesothelioma deaths on the mesothelioma register for the period 2001-2021. This is mainly due to the age restriction of 16-74 years (the age range for which last occupation of the deceased is routinely recorded on death certificates in England and Wales), but also due to missing or invalid occupation codes for some deaths below age 75 years (1.5% of male and 9.2% of female deaths).

Death data for all causes of death combined required for the calculation of PMRs from 2011-20 for England and Wales were supplied by the Office for National Statistics (ONS). For deaths registered after 1 April 2011, occupations have been classified according to the Standard Occupational Classification 2010 (SOC2010), and for deaths during 2001-2010 occupations have been classified according to the Standard Occupational Classification 2000 (SOC2000).

A small number of deaths in this analysis occurring during 2001-2010 originally coded to either SOC90 or SOC2010 were recoded SOC2000 using a probability matching algorithm provided by the ONS combined with additional checks made against the job description.

Information about the Standard Occupational Classification is available from the Office for National Statistics:

www.ons.gov.uk/methodology/classificationsandstandards/standardoccupationalclassification/soc

In this analysis, mortality in the different occupational groups is represented by Proportional Mortality Ratios (PMRs) and associated 95% confidence intervals. A PMR for a particular occupation is the ratio of the observed number of deaths for that occupation to the expected number of deaths, with that ratio expressed as a percentage (i.e. multiplied by 100).

The expected number of deaths is calculated as the number of mesothelioma deaths that would have been recorded for that occupation if the proportion of mesothelioma deaths was equal to the proportion of total deaths from all causes in that occupation. Since mesothelioma incidence is also strongly related to age, the calculation also takes account of differences in the distribution of ages between occupational groups. A worked example of how the PMR is calculated for a particular occupation is given below.

Statistics have been calculated for 1 to 4 digit codes i.e. major, sub-major, minor, and unit groups of SOC2010 for the period 2011-20 and SOC2000 for the period 2001-10.

The statistical models shown in the graphs, involved fitting a smoothed term for the year in a Poisson Generalized Additive model (GAM) to identify annual trends. In a most cases a Poisson error term was assumed; for a small number of cases a Negative Binomial or Normal (Gaussian) error term was assumed as this provided a better fit to the data.

Example PMR calculation

The table below illustrates the calculation of a PMR for men in “occupation X”. Column 3 gives the proportion of all mesothelioma deaths by age (=column 2 divided by column 1). This proportion is applied to the number of deaths from all causes by age in occupation X, given in column 4, to give the expected number of deaths from mesothelioma in this occupation in column 5. The total observed number of mesothelioma deaths in occupation X was 500 (not shown in table). Dividing this by the total expected number of deaths (sum of column 5 = 230 deaths) expressed as a percentage gives a PMR of 217 in this case.

Age group	Deaths				
	All men			Men in occupation X	
	All causes (1)	Mesothelioma deaths (2)	Proportion from mesothelioma (3) = (2) / (1)	All-cause deaths (4)	Expected deaths (5) = (3) * (4)
16-19	16,500	1	0.000061	6,400	0.388
20-24	21,732	1	0.000046	7,833	0.360
25-29	18,072	5	0.000277	7,907	2.188
30-34	20,544	16	0.000779	7,770	6.051
35-39	27,300	76	0.002784	6,443	17.937
40-44	42,576	199	0.004674	6,222	29.082
45-49	61,236	402	0.006565	6,243	40.984
50-54	102,900	705	0.006851	6,391	43.787
55-59	187,416	1,145	0.006109	6,269	38.300
60-64	308,988	1,436	0.004647	5,367	24.943
65-69	433,956	1,499	0.003454	4,997	17.261
70-74	550,296	1,315	0.002390	3,729	8.911
All ages 16-74	1,791,516	6,800		75,571	230

Confidence intervals and statistical significance

A PMR calculated for an occupational group may be greater or less than 100 by chance. Confidence intervals are used to give an indication of the uncertainty associated with each PMR due to this random variation. A 95% confidence interval is such that, if the calculation could be repeated many times with different samples of the events, then the confidence interval will contain the true value of the PMR 95% of the time. If the lower confidence limit is greater than 100 then the PMR is said to be statistically significantly elevated. Likewise, if the upper confidence interval that is presented is lower than 100 then the PMR is said to be statistically significantly reduced. In this analysis, confidence intervals are calculated assuming Poisson variability in the mesothelioma count for each occupation.

National Statistics

National Statistics status means that statistics meet the highest standards of trustworthiness, quality and public value. They are produced in compliance with the Code of Practice for Statistics, and awarded National Statistics status following assessment and compliance checks by the Office for Statistics Regulation (OSR). The last compliance check of these statistics was in 2013.

It is Health and Safety Executive's responsibility to maintain compliance with the standards expected by National Statistics. If we become concerned about whether these statistics are still meeting the appropriate standards, we will discuss any concerns with the OSR promptly. National Statistics status can be removed at any point when the highest standards are not maintained, and reinstated when standards are restored.

Details of OSR reviews undertaken on these statistics, quality improvements, and other information noting revisions, interpretation, user consultation and use of these statistics is available from www.hse.gov.uk/statistics/about.htm

An account of how the figures are used for statistical purposes can be found at www.hse.gov.uk/statistics/sources.htm.

For information regarding the quality guidelines used for statistics within HSE see www.hse.gov.uk/statistics/about/quality-guidelines.htm

A revisions policy and log can be seen at www.hse.gov.uk/statistics/about/revisions/

Additional data tables can be found at www.hse.gov.uk/statistics/tables/.

General enquiries: Statistician: Lucy.Darnton@hse.gov.uk

Journalists/media enquiries only: www.hse.gov.uk/contact/contact.htm





Further information

For information about health and safety, or to report inconsistencies or inaccuracies in this guidance, visit [the HSE website](#).

You can order HSE priced publications at [the HSE books website](#).

HSE priced publications are also available from bookshops.

This publication is available on the HSE website www.hse.gov.uk/statistics/causdis

© Crown copyright If you wish to reuse this information visit [the HSE website](#) for details. First published 07/22.

Published by the Health and Safety Executive 07/22.