

Mesothelioma deaths by Geographical Area, 2026



Mesothelioma deaths for Local and Unitary Authority areas in Great Britain 1981-2024

July 2026



Contents

Summary	2
Introduction	3
Results and Discussion	6
Results for the overall period 1981-2024	7
Temporal trends – Scotland, Wales and English regions	15
Temporal trends for UA and LA areas, males	17
Temporal trends for UA and LA areas, females	20
Annex 1 – Methodology	24
SMR calculation – worked example	24
Annex 2 – Unitary and Local Authorities as of 2015	26
Annex 3 – Other areas with significantly raised male SMRs	28
References	32
Accredited Official Statistics	33

Summary

The information in this document relates to Health and Safety Statistics published by the Health and Safety Executive in 2026. The document can be found at:

www.hse.gov.uk/statistics/causdis

This fact sheet provides statistics on mesothelioma deaths in Great Britain by geographical area for deaths occurring in the period 1981 to 2024. Numbers of deaths for males and females are given for areas within the current local government structure down to Unitary Authority (UA) and Local Authority (LA) level.

Standardised Mortality Ratios (SMRs) are also provided with associated 95% Confidence Intervals to allow comparison of areas after accounting for differences in the age distributions of the underlying populations.

The statistics are presented in tabular form in the following spreadsheet:

www.hse.gov.uk/statistics/assets/docs/mesoarea.xlsx

- MESOAREA01: Number of mesothelioma deaths and SMRs for males by geographical area in Great Britain.
- MESOAREA02: Number of mesothelioma deaths and SMRs for females by geographical area in Great Britain.
- MESOAREA03: Mesothelioma deaths and Standardised Mortality Ratios (SMRs) for males in Great Britain by area and five-year time periods 1981-2024.
- MESOAREA04: Mesothelioma deaths and Standardised Mortality Ratios (SMRs) for females in Great Britain by area and five-year time periods 1981-2024.

Results are also available as interactive maps at: <https://arcg.is/1qO0G40>.

Introduction

Previous descriptive analyses of mesothelioma death rates for geographical areas within Great Britain have highlighted the effect of geographically-specific sources of asbestos exposure [1]. Geographical areas with the highest mortality rates tend to be those known to contain large industrial sites such as shipyards and asbestos product factories.

This factsheet provides an update of analyses of mesothelioma mortality by Unitary Authority (UA) and Local Authority (LA) area to include deaths occurring during the period 1981 to 2024, the longest period for which data are available according to the current UA and LA structure. It also provides more detailed statistical modelling of temporal trends within these geographical areas (see Annex 1 for further details).

The analyses are based on the last area of residence of the deceased, as recorded on death certificates, and use Standardised Mortality Ratios (SMRs) which compare the mortality rate in a particular area with the mortality rate for GB, taking account of age differences (see Annex 1 for further details). SMRs are expressed as a percentage: values higher or lower than 100 indicate mesothelioma rates that are higher or lower, respectively, than for GB as a whole.

A number of Unitary Authorities were created in 2009 and these are detailed in Annex 2, along with changes in 2018, 2019, 2020, 2021 and 2023.

The analyses of temporal trends for geographical areas within Great Britain should be interpreted in the context of increasing annual mesothelioma deaths in Great Britain as a whole over the period of the analysis. There were nearly 4 times as many deaths in the latest decade (2015-24) compared with the 1980s (Table 1).


Table 1: Male and female mesothelioma deaths 1981-2024(p) by time period.

Time period	Male deaths	Female deaths	Total deaths
1981-1994	9,906	1,586	11,492
1995-2004	14,064	2,314	16,378
2005-2014	19,428	3,748	23,176
2015-2024	19,922	4,183	24,105

Similar patterns in annual numbers over time are evident for both males and females, though annual male deaths have consistently outnumbered female deaths by around five to one due to higher and more widespread past asbestos exposures often in occupational settings (Figure 1).

Since Standardised Mortality Ratios (SMRs) compare the mortality rate in a particular region with that for GB as a whole, trends in SMRs for a particular area indicate whether rates for that area have increased relatively more or less rapidly than for GB as a whole. No change in the SMR for an area over time indicates that the mortality rates have increased in line with the trend for GB as a whole.

Figure 1: Male and female mesothelioma deaths 1968-2024(p)



Figures for 2024 are provisional.

All of the analyses presented in this factsheet are limited by the fact that death certificates record only the last address of residence of the deceased. A case of mesothelioma caused by work in one geographical area will only be assigned to that area in this analysis if the individual was resident there when they died. The long latency period of mesothelioma means that individuals may move between areas before the onset of the disease and thus there is considerable potential for dilution of the observed difference in risk between areas. The extent of this dilution will be strongest for those areas where there have been substantial migrations. Areas with the highest SMRs will be those which are genuine sources of risk, but their SMRs will understate the true risk level relative to the rest of Great Britain. Conversely, SMRs of other areas will overstate the level of risk associated with these locations. The areas recording the lowest SMRs will be those areas not

associated with asbestos exposure and which are unlikely to be the final area of residence for individuals with asbestos exposure.

Results and Discussion

These analyses for the period 1981 to 2024 are based on 63,320 male and 11,831 female mesothelioma deaths (a small number of individuals with an overseas address are excluded).

Full results are available in Excel tables at:

www.hse.gov.uk/statistics/assets/docs/mesoarea.xlsx

- MESOAREA01: Number of mesothelioma deaths and SMRs for males by geographical area in Great Britain.
- MESOAREA02: Number of mesothelioma deaths and SMRs for females by geographical area in Great Britain.
- MESOAREA03: Mesothelioma deaths and Standardised Mortality Ratios (SMRs) for males in Great Britain by area and five-year time periods 1981-2024.
- MESOAREA04: Mesothelioma deaths and Standardised Mortality Ratios (SMRs) for females in Great Britain by area and five-year time periods 1981-2024.

Maps showing SMRs for males and females for the overall period 1981-2024 are presented (Figures 2 and 4) along with additional maps highlighting those areas for which the mortality rate was statistically significantly higher or lower than for GB as a whole (Figures 3 and 5).

Results are also available as interactive maps available at: <https://arcg.is/1qO0G40>.

Temporal trends for Scotland, Wales and English regions are shown for males and females in Figures 6 and 7, and trends for selected LA and UA areas with higher SMRs are shown in Figures 8-12 for males and Figures 13-17b for females. Additional results for males are available and in Annex 3 (Figures 18-22).

Where results are presented in the text that follows, we provide the SMR followed by its 95% Confidence Interval in brackets and the number of observed deaths that the SMR is based on.

Results for the overall period 1981-2024

The geographical areas with the highest male mesothelioma death rates for the period 1981-2024 were:

- 1. West Dunbartonshire: SMR 337.1 (300.5 to 377), n=308
- 2. South Tyneside: SMR 272.5 (248.4 to 298.3), n=468
- 3. North Tyneside: SMR 271.9 (250.6 to 294.5), n=604
- 4. Portsmouth: SMR 256.5 (234.1 to 280.5), n=482
- 5. Plymouth: SMR 256.2 (237.3 to 276.3), n=674
- 6. Medway: SMR 228.8 (209.7 to 249.2), n=527
- 7. Hartlepool: SMR 215.1 (186.8 to 246.5), n=207
- 8. Barking and Dagenham: SMR 212.1 (188.9 to 237.4), n=304
- 9. Southampton: SMR 211.1 (191.8 to 231.8), n=439
- 10. Gosport: SMR 209.9 (180.1 to 243.3), n=177

The geographical areas with the highest female mesothelioma death rates were:

- 1. Barking and Dagenham: SMR 313.3 (252.3 to 384.7), n=91
- 2. Sunderland: SMR 302.6 (258.8 to 351.7), n=170
- 3. Newham: SMR 268.1 (213.2 to 332.8), n=82
- 4. West Dunbartonshire: SMR 219.9 (158.5 to 297.2), n=42
- 5. Leeds: SMR 219.8 (195.9 to 245.9), n=306
- 6. Castle Point: SMR 217.3 (156.6 to 293.7), n=42
- 7. Havering: SMR 210.5 (171.8 to 255.3), n=103
- 8. Basildon: SMR 199 (152.2 to 255.6), n=61
- 9. Medway: SMR 189.4 (150.5 to 235.5), n=81
- 10. Rochford: SMR 187.8 (129.3 to 263.7), n=33

Due to local government changes in 2023, Barrow-in-Furness – the area that had previously had the highest male SMR (399.3) and the 6th highest female SMR (213.9) – became part of the new Westmorland and Furness unitary authority.

As in previous geographical analyses of mesothelioma deaths, the results presented here show that areas with the highest excess of mesothelioma in males tend to be those containing industrial sites known to have been associated with high asbestos exposures in the past, such as shipyards. However, other evidence suggest that asbestos exposures in the construction industry account for a substantial proportion of mesothelioma deaths (for example an epidemiological study in GB estimated that 46% of mesotheliomas among men born in the 1940s were attributed to such exposures). Such exposures are less likely

to have been associated with specific geographical areas; rather, they are likely to have taken place over a wide range of areas.

The analyses of temporal trends show that most areas associated with shipbuilding activity tend to have much higher SMRs for early time periods than for later periods, although there are some exceptions. Mesothelioma rates in these areas thus tend to have risen more slowly over time than the overall rate for Great Britain. This may to some extent reflect the effect of risks being diluted due to the migration of those exposed in an industry which has declined substantially into lower risk areas. It may also suggest that annual mesothelioma deaths arising from such exposures may have peaked earlier than those arising from other sources of exposure.




Figure 2 – Mesothelioma SMRs for males by unitary/local authority for the period 1981-2024




Figure 3 – Statistical significance of mesothelioma SMRs for males by unitary/local authority 1981-2024




Figure 3A – Mesothelioma SMRs for males by unitary/local authority by 10-year periods 1981-2024




Figure 4 – Mesothelioma SMRs for females by unitary/local authority for the period 1981-2024




Figure 5 – Statistical significance of mesothelioma SMRs for females by unitary/local authority 1981-2024




Figure 5A – Mesothelioma SMRs for females by unitary/local authority by 10-year period 1981-2024

Temporal trends – Scotland, Wales and English regions

Figure 6 shows the regional variation for male SMRs calculated annually along with 95% confidence intervals. Corresponding statistics for females are shown in Figure 7.

Trend lines with solid bold **black** lines indicate a statistically significant yearly trend, those with **green** lines indicate trends of borderline significance, and for those with **blue** lines trends were not significant. The dashed lines represent the 95% confidence intervals.




Figure 6 – Annual mesothelioma SMRs for males by region, 1981-2024

For males, the highest rates for the period as a whole (1981-2024) were seen in the North East, South East and East of England. SMRs have declined over time in the North East, Scotland and London. This indicates that mesothelioma rates for these regions have increased relatively less rapidly over the period than for GB as a whole.

Conversely, there was an increase in the SMRs over time for those regions with the lowest rates for the period as a whole (East Midlands, West Midlands and Wales). This indicates that mesothelioma rates for these regions have increased relatively more rapidly over the period than for GB as a whole.

England accounts for the majority of Great Britain and so SMRs show little variation from the standard SMR figure of 100.



Figure 7 – Annual mesothelioma SMRs for females by region, 1981-2024

For females, the highest rates for the period as a whole were in the North-East, East followed by South East, Yorkshire and The Humber and London. There was evidence of an increase in the SMR over time for the East, South East, South West and Wales, with evidence of a decline for the North East and London.

Temporal trends for UA and LA areas, males

Temporal trends in annual male SMRs are shown in Figures 8-12 (and Figures 18-21 in Annex 3) for all UA/LA areas with significantly elevated SMRs for the period as a whole (1981-2024).

Figure 8 covers the top six areas all of which were associated with shipbuilding.

Figure 9 covers the North East, Teesside and Tyne and Wear (North Tyneside and South Tyneside are included in Figure 8).

Figure 10 covers the coastal region of Hampshire (except for Portsmouth which is included in Figure 8).

Figure 11 covers parts of central Scotland, mainly around the Clyde (except West Dunbartonshire which is included in Figure 8).




Figure 8 – Annual mesothelioma SMRs for males for the top six UA/LA areas, 1981-2024

Marked statistically significant downward trends were seen for these areas.

Prior to the local government reorganisation, Barrow-in-Furness had consistently had the highest SMR, although this reduced from over 900 to less than 300 over the period, with an SMR of 399.3 for the period 1981-2022).

For Renfrewshire (Figure 11) a decline was seen in the SMR from around 350 in 1981 to being not statistically different from 100 in 2024.




Figure 9 – Annual mesothelioma SMRs for males for UA/LAs in the North East, 1981-2024




Figure 10 – Annual mesothelioma SMRs for males for UA/LAs Hampshire coast 1981-2024




Figure 11 – Annual mesothelioma SMRs for males for UA/LAs in central Scotland, 1981-2024




Figure 12 – Annual mesothelioma SMRs for males for other areas, 1981-2024

Several areas with high SMRs for males also had high SMRs for females including Sunderland, Barking and Dagenham, and West Dunbartonshire.

Temporal trends for UA and LA areas, females

Temporal trends in annual female SMRs are shown in Figures 13-17 for all areas with significantly elevated SMRs for the period as a whole (1981-2024).

Figure 13 shows marked statistically significant declines in the annual SMRs for Barking and Dagenham, Sunderland. SMRs for Blackburn with Darwen were not significantly raised in more recent years.

Mesothelioma mortality statistics by geographical area for Great Britain, 2026




Figure 13 – Annual mesothelioma SMRs for females for the top eight UA/LA areas, 1981-2024




Figure 14 – Annual mesothelioma SMRs for females for areas ranked 9-16 in GB, 1981-2024

Mesothelioma mortality statistics by geographical area for Great Britain, 2026




Figure 15 – Annual mesothelioma SMRs for females for areas ranked 17-24 in GB, 1981-2024




Figure 16 – Annual mesothelioma SMRs for females for other areas, 1981-2024




Figure 17 – Annual mesothelioma SMRs for females for other areas, 1981-2024 (part 2)

Annex 1 – Methodology

Mesothelioma deaths occurring during the period 1981-2024 were obtained from the Health and Safety Executive Mesothelioma Register. SMRs were derived using mid-year population estimates provided by the Office for National Statistics.

The method of age standardisation used in the production of SMRs is commonly referred to as the indirect method. Age-specific death rates in a standard population (in this case Great Britain by gender) are applied to the age structure of the population for each geographical area in order to calculate expected numbers of deaths. The ratio of the observed number of deaths to the expected number of deaths in the area is calculated and multiplied by 100 to give the SMR. The SMR of the standard population is 100. An SMR greater or less than 100 indicates a respectively higher or lower than expected mortality rate in a specific area. If the lower bound of the 95% Confidence Interval for the SMR is greater than 100 this indicates that the observed number of deaths was statistically significantly higher than expected. A worked example of the SMR calculation is provided below.

The statistical models used for the analysis of time trends in this report are Generalized Additive Models (GAMs). These models permit a flexible analysis of trends which do not depend on assuming a particular mathematical relationship in advance, such as a linear decline over the entire time period. Instead, the GAMs allow the fitting of trends which take into account different patterns of change within different parts of the overall time period and smooth out some of the year-on-year variation. In most cases the models assume a Poisson error term. In a small number of cases a Negative Binomial or Normal (Gaussian) error term was assumed.

SMR calculation – worked example

Table A illustrates the calculation of an SMR for men in area A. The total population of Great Britain is used as the standard population, (column 1). The mesothelioma death rate per 1,000 in the population for each age group (column 3) is the total number of male mesothelioma deaths (column 2) divided by the total number of men in Great Britain (column 1) to give age-specific death rates in the standard population. These rates are applied to the total population in area A, given in column 4, to give the expected numbers of deaths in this area, in column 6. The total observed number of deaths, shown in column 5, (1,196) divided by the expected number of deaths (2,024), multiplied by 100 gives an SMR of 59.

Age group	All men			Men in area A		
	Population, (thousands)	Mesothelioma deaths	Death rate Per 1,000	Population, (thousands)	Mesothelioma deaths	Expected deaths
	(1)	(2)	(3) = (2) / (1)	(4)	(5)	(6)=(3)*(4)
0-4	28 554	0	0.000	6 926	0	0.0
5-9	29 683	0	0.000	8 514	0	0.0
10-14	32 324	0	0.000	9 286	0	0.0
15-19	35 061	1	0.000	8 729	0	0.2
20-24	34 931	1	0.000	7 833	0	0.2
25-29	32 949	5	0.000	7 907	2	1.2
30-34	31 188	16	0.001	7 770	7	3.9
35-39	29 220	76	0.003	6 443	17	16.7
40-44	27 454	199	0.007	6 222	32	45.1
45-49	24 983	402	0.016	6 243	76	100.4
50-54	24 398	705	0.029	6 391	136	184.7
55-59	24 001	1 145	0.048	6 269	179	299.1
60-64	22 155	1 436	0.065	5 367	183	347.9
65-69	19 554	1 499	0.077	4 997	222	383.1
70-74	15 232	1 315	0.086	3 729	177	321.9
75-79	10 232	930	0.091	2 176	112	197.8
80-84	5 176	472	0.091	1 007	40	91.8
85+	2 503	145	0.058	525	13	30.4
All ages	429 600	8 347		106 334	1 196	2024.5

Table A: Example of an SMR calculation

Annex 2 – Unitary and Local Authorities as of 2015

The following map shows all UAs and LAs as of 2015:

data.gov.uk/dataset/f96d4afa-1100-484d-9f38-abf79a830db7/local-authority-districts-counties-and-unitary-authorities-december-2015-map-in-great-britain.

This structure comprises the following local authorities that existed as part of the previous local government structure that applied from 1998. Changes in 2009 were as follows:

1. County Durham UA comprises the former districts of Chester-le-Street, Derwentside, Durham, Easington, Sedgefield, Teesdale and Wear Valley.
2. Northumberland UA comprises the former districts of Alnwick, Berwick-upon-Tweed, Blyth Valley, Castle Morpeth, Tynedale and Wansbeck.
3. Cheshire East UA comprises the former districts of Congleton, Crewe and Nantwich and Macclesfield.
4. Cheshire West and Chester UA comprises the former districts of Chester, Ellesmere Port & Neston and Vale Royal.
5. Shropshire UA comprises the former districts of Bridgnorth, North Shropshire, Oswestry, Shrewsbury and Atcham and South Shropshire.
6. Bedford UA comprises the former district of Bedford.
7. Central Bedfordshire UA comprises the former districts of Mid Bedfordshire and South Bedfordshire.
8. Cornwall UA comprises the former districts of Caradon, Carrick, Kerrier, North Cornwall, Penwith and Restormel.
9. Wiltshire UA comprises the former districts of Kennet, North Wiltshire, Salisbury and West Wiltshire.

More recent changes include:

10. Tyne and Wear (Met County) is now code E11000007, it was previously E11000004
11. In 2018 Fife changed from S12000015 to S12000047, and Perth and Kinross from S12000024 to S12000048 due to a boundary change.

12. In 2019 Glasgow City changed from S12000046 to S12000049, and North Lanarkshire from S12000044 to S12000050 due to a boundary change.

13. In 2019 the new unitary authority of Bournemouth, Christchurch and Poole (E06000058) was created from Bournemouth (E06000028), Poole (E06000029) and Christchurch (E07000048); along with the new unitary authority of Dorset (E06000059) replacing the old county minus Christchurch i.e. for East Dorset (E07000049), North Dorset (E07000050), Purbeck (E07000051), West Dorset (E07000052) and Weymouth and Portland (E07000053). New district councils were created for Somerset West and Taunton (E07000246) from Taunton Deane (E07000190) and West Somerset (E07000191); West Suffolk (E07000245) from Forest Heath (E07000201) and St Edmundsbury (E07000204); East Suffolk (E07000244) from Suffolk Coastal (E07000205) and Waveney (E07000206).

14. In 2020 Buckinghamshire unitary authority (E06000060) replaced the county (E10000002) and the 4 local authorities of Aylesbury Vale (E07000004), Chiltern (E07000005), South Bucks (E07000006) and Wycombe (E07000007).

15. On 1 April 2021, the non-metropolitan county of Northamptonshire (E10000021) and its seven districts were abolished, and two new unitary authorities were created:

North Northamptonshire (E06000061), consisting of the old non-metropolitan districts of Corby (E07000150), East Northamptonshire (E07000152), Kettering (E07000153), and Wellingborough (E07000156).

West Northamptonshire (E06000062), consisting of the old non-metropolitan districts of Daventry (E07000151), Northampton (E07000154), and South Northamptonshire (E07000155).

16. On 1st April 2023 the following new unitary authorities were created:

County of Cumbria (E10000006) and its districts were replaced by 2 unitary authorities; Cumberland (E06000063) replaced Allerdale (E07000026), Carlisle (E07000028) and Copeland (E07000029); and Westmorland and Furness (E06000064) replaced Barrow-in-Furness (E07000027), Eden (E07000030) and South Lakeland (E07000031).

The county of North Yorkshire (E10000023) and its districts were replaced by a new North Yorkshire (E06000065) unitary authority, the districts being Craven (E07000163), Hambleton (E07000164), Harrogate (E07000165), Richmondshire (E07000166), Ryedale (E07000167), Scarborough (E07000168) and Selby (E07000169).

Somerset (E06000066) replaced the county of Somerset (E07000246) and its districts Mendip (E07000187), Sedgemoor (E07000188), South Somerset (E07000189) and Somerset West and Taunton (E07000246).

Annex 3 – Other areas with significantly raised male SMRs




Figure 18 – Annual mesothelioma SMRs for males for other areas, 1981-2024 (part 2)




Figure 19 – Annual mesothelioma SMRs for males for other areas, 1981-2024 (part 3)

Mesothelioma mortality statistics by geographical area for Great Britain, 2026




Figure 20 – Annual mesothelioma SMRs for males for other areas, 1981-2021 (part 4)

Mesothelioma mortality statistics by geographical area for Great Britain, 2026




Figure 21 – Annual mesothelioma SMRs for males for other areas, 1981-2024 (part 5)

References

1. McElvenny D, Darnton A, Price M, Hodgson J. Mesothelioma mortality in Great Britain from 1968 to 2001. *Occup Med* 2005;55(2):79-87.

Accredited Official Statistics

National Statistics are accredited official statistics. This publication is part of HSE's accredited official statistics releases. <https://uksa.statisticsauthority.gov.uk/about-the-authority/uk-statistical-system/types-of-official-statistics/>

HSE's official statistics practice is regulated by the Office for Statistics Regulation (OSR). Accredited Official Statistics are a subset of official statistics that have been independently reviewed by the OSR and confirmed to comply with the standards of trustworthiness, quality and value in the Code of Practice for Statistics. Accredited official statistics were previously called National Statistics (and still referenced as such in Statistics and Registration Service Act 2007). See uksa.statisticsauthority.gov.uk/about-the-authority/uk-statistical-system/types-of-official-statistics/ for more details on the types of official statistics.

From 7 June 2024 the Accredited Official Statistics badge has replaced the previous National Statistics badge. These statistics were last reviewed by OSR in 2013. It is Health and Safety Executive's responsibility to maintain compliance with the standards expected. If we become concerned about whether these statistics are still meeting the appropriate standards, we will discuss any concerns with the OSR promptly. Accredited Official Statistics status can be removed at any point when the highest standards are not maintained, and reinstated when standards are restored.

Details of OSR reviews undertaken on these statistics, quality improvements, and other information noting revisions, interpretation, user consultation and use of these statistics is available from www.hse.gov.uk/statistics/about.htm.

An account of how the figures are used for statistical purposes can be found at www.hse.gov.uk/statistics/sources.htm.

For information regarding the quality guidelines used for statistics within HSE see www.hse.gov.uk/statistics/about/quality-guidelines.htm

A revisions policy and log can be seen at www.hse.gov.uk/statistics/about/revisions/

Additional data tables can be found at www.hse.gov.uk/statistics/tables/.

Lead Statistician: Kevin Shepherd

Feedback on the content, relevance, accessibility and timeliness of these statistics and any non-media enquiries should be directed to:

Email: statsfeedback@hse.gov.uk

Journalists/media enquiries only: www.hse.gov.uk/contact/contact.htm





Further information

For information about health and safety, or to report inconsistencies or inaccuracies in this guidance, visit [the HSE website](#).

You can order HSE priced publications at [the HSE books website](#).

HSE priced publications are also available from bookshops.

This publication is available on the HSE website www.hse.gov.uk/statistics/causdis

© Crown copyright If you wish to reuse this information visit [the HSE website](#) for details. First published 07/26.

Published by the Health and Safety Executive 07/26.